


VideoMot2

References



Publications VideoMot2

2015

- **Cathomas F, Stegen M, Sigrist H, Schmid L, Seifritz E, Gassmann M, Bettler B, Pryce CR.** Altered emotionality and neuronal excitability in mice lacking KCTD12, an auxiliary subunit of GABAB receptors associated with mood disorders. *Transl Psychiatry* 2015; 5:e510
<http://www.ncbi.nlm.nih.gov/pubmed/25689571>
- **Jain P, Hassan AM, Koyani CN, Mayerhofer R, Reichmann F, Farzi A, Schuligoi R, Malle E, Holzer P.** Behavioral and molecular processing of visceral pain in the brain of mice: impact of colitis and psychological stress.
<http://www.ncbi.nlm.nih.gov/pubmed/26217204>
- **Piao J, Major T, Auyeung G, Policarpio E, Menon J, Droms L, Gutin P, Uryu K, Tchieu J, Soulet D, Tabar V.** Human embryonic stem cell-derived oligodendrocyte progenitors remyelinate the brain and rescue behavioral deficits following radiation. *Cell Stem Cell* 2015; 16(2):198-210
<http://www.ncbi.nlm.nih.gov/pubmed/25658373>
- **Reichmann F, Painsipp E, Holzer P, Kummer D, Bock E, Leitinger G.** A novel unbiased counting method for the quantification of synapses in the mouse brain. *J Neurosci Methods* 2015; 240:13-21
<http://www.ncbi.nlm.nih.gov/pubmed/25445248>

2014

- **Amendola E, Zhan Y, Mattucci C, Castroflorio E, Calcagno E, Fuchs C, Lonetti G, Silingardi D, Vysotski AL, Farley D, Ciani E, Pizzorusso T, Giustetto M, Gross CT.** Mapping pathological phenotypes in a mouse model of CDKL5 disorder. *PLoS One* 2014; 9(5): e91613
<http://www.ncbi.nlm.nih.gov/pubmed/24838000>
- **Anier K, Malinovskaja K, Pruus K, Aonurm-Helm A, Zharkovskiy A, Kalda A.** Maternal separation is associated with DNA methylation and behavioural changes in adult rats. *Eur Neuropsychopharmacol* 2014; 24(3): 459-68
<http://www.ncbi.nlm.nih.gov/pubmed/23972903>
- **An XL, Tai FD.** AVP and Glu systems interact to regulate levels of anxiety in BALB/cJ mice. *Zool Res* 2014; 35(4): 319-25
<http://www.ncbi.nlm.nih.gov/pubmed/25017752>
- **Balseanu AT, Buga AM, Catalin B, Wagner DC, Boltze J, Zagrean AM, Reymann K, Schaebitz W, Popa-Wagner A.** Multimodal approaches for regenerative stroke therapies: combination of granulocyte colony-stimulating factor with bone marrow mesenchymal stem cells is not superior to G-CSF alone. *Front Aging Neurosci* 2014; 6:130
<http://www.ncbi.nlm.nih.gov/pubmed/25002846>
- **Bhattacharya S, Haertel C, Maelicke A, Montag D.** Galantamine slows down plaque formation and behavioral decline in the 5XFAD mouse model of Alzheimer's disease. *PLoS One* 2014; 9(2): e89454
<http://www.ncbi.nlm.nih.gov/pubmed/24586789>

- **Brunner SM, Farzi A, Locker F, Holub BS, Drexel M, Reichmann F, Lang AA, Mayr JA, Vilches JJ, Navarro X, Lang R, Sperk G, Holzer P, Kofler B.** GAL3 receptor KO mice exhibit an anxiety-like phenotype. *Proc Natl Acad Sci USA* 2014; 111(19): 7138-43
<http://www.ncbi.nlm.nih.gov/pubmed/24782539>
- **Byrns CN, Pitts MW, Gilman CA, Hashimoto AC, Berry MJ.** Mice lacking selenoprotein P and selenocysteine lyase exhibit severe neurological dysfunction, neurodegeneration, and audiogenic seizures. *J Biol Chem* 2014; 298(14):9662-74
<http://www.ncbi.nlm.nih.gov/pubmed/24519931>
- **Carola V, Perlas E, Zonfrillo F, Soini HA, Novotny MV, Gross CT.** Modulation of social behavior by the agouti pigmentation gene. *Front Behav Neurosci* 2014; 8:259
<http://www.ncbi.nlm.nih.gov/pubmed/25136298>
- **Farzi A, Reichmann F, Meinitzer A, Mayerhofer R, Jain P, Hassan AM, Fröhlich EE, Wagner K, Painsipp E, Rinner B, Holzer P.** Synergistic effects of NOD1 or NOD2 and TLR4 activation on mouse sickness behavior in relation to immune and brain activity markers. *Brain Behav Immun* 2014; Epub ahead of print
<http://www.ncbi.nlm.nih.gov/pubmed/25218901>
- **Frey A, Popp S, Post A, Langer S, Lehmann M, Hofmann U, Sirén AL, Hommers L, Schmitt A, Strekalova T, Ertl G, Lesch KP, Frantz S.** Experimental heart failure causes depression-like behavior together with differential regulation of inflammatory and structural genes in the brain. *Front Behav Neurosci* 2014; 8:376
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4215623>
- **Grieb B, Engler G, Sharott A, von Nicolai C, Streichert T, Papageorgiou I, Schulte A, Westphal M, Lamszus K, Engel AK, Moll CK, Hamel W.** High-frequency stimulation of the subthalamic nucleus counteracts cortical expression of major histocompatibility complex genes in a rat model of Parkinson's disease. *PLoS One* 2014; 9(3):e91663
<http://www.ncbi.nlm.nih.gov/pubmed/24621597>
- **Hassan AM, Jain P, Reichmann F, Mayerhofer R, Farzi A, Schuligoi R, Holzer P.** Repeated predictable stress causes resilience against colitis-induced behavioral changes in mice. *Front Behav Neurosci* 2014; 8:386
<http://www.ncbi.nlm.nih.gov/pubmed/25414650>
- **Issler O, Carter RN, Paul ED, Kelly PA, Olverman HJ, Neufeld-Cohen A, Kuperman Y, Lowry CA, Seckl JR, Chen A, Jamieson PM.** Increased anxiety in corticotropin-releasing factor type 2 receptor-null mice requires recent acute stress exposure and is associated with dysregulated serotonergic activity in limbic brain areas. *Biol Mood Anxiety Disord* 2014; 4(1):1
<http://www.ncbi.nlm.nih.gov/pubmed/24447313>
- **Issler O, Haramati S, Paul ED, Maeno H, Navon I, Zwang R, Gil S, Mayberg HS, Dunlop BW, Menke A, Awatramani R, Binder EB, Deneris ES, Lowry CA, Chen A.** MicroRNA 135 is essential for chronic stress resiliency, antidepressant efficacy, and intact serotonergic activity. *Neuron* 2014; 83(2):344-60
<http://www.ncbi.nlm.nih.gov/pubmed/24952960>
- **Mihailovska E, Raith M, Valencia RG, Fischer I, Al Banchaabouchi M, Herbst R, Wiche G.** Neuromuscular synapse integrity requires linkage of acetylcholine receptors to postsynaptic intermediate filament networks via rapsyn-plectin 1f complexes. *Mol Biol Cell* 2014; 25(25):4130-49
<http://www.ncbi.nlm.nih.gov/pubmed/25318670>

- **Örd T, Innos J, Lilleväli K, Tekko T, Sütt S, Örd D, Köks S, Vasar E, Örd T.** Trib3 is developmentally and nutritionally regulated in the brain but is dispensable for spatial memory, fear conditioning and sensing of amino acid-imbalanced diet. *PLoS One* 2014; 9(4):e94691
<http://www.ncbi.nlm.nih.gov/pubmed/24732777>
- **Seo J, Giusti-Rodríguez P, Zhou Y, Rudenko A, Cho S, Ota KT, Park C, Patzke H, Madabhushi R, Pan L, Mungenast AE, Guan JS, Delalle I, Tsai LH.** Activity-dependent p25 generation regulates synaptic plasticity and A β -induced cognitive impairment. *Cell* 2014; 157(2):486-98
<http://www.ncbi.nlm.nih.gov/pubmed/24725413>
- **Tauber SC, Harms K, Falkenburger B, Weis J, Sellhaus B, Nau R, Schulz JB, Reich A.** Modulation of hippocampal neuroplasticity by Fas/CD95 regulatory protein 2 (Faim2) in the course of bacterial meningitis. *J Neuropathol Exp Neurol* 2014; 73(1):2-13
<http://www.ncbi.nlm.nih.gov/pubmed/24335530>
- **Volk N, Paul ED, Haramati S, Eitan C, Fields BK, Zwang R, Gil S, Lowry CA, Chen A.** MicroRNA-19b associates with Ago2 in the amygdala following chronic stress and regulates the adrenergic receptor beta 1. *J Neurosci* 2014; 34(45):15070-82
<http://www.ncbi.nlm.nih.gov/pubmed/25378171>
- **Wegener E, Brendel C, Fischer A, Hülsmann S, Gärtner J, Huppke P.** Characterization of the MeCP2R168X Knockin Mouse Model for Rett Syndrome. *PLoS One* 2014; 9(12): e115444
<http://www.ncbi.nlm.nih.gov/pubmed/25541993>
- **Zhan Y, Paolicelli RC, Sforzini F, Weinhard L, Bolasco G, Pagani F, Vyssotski AL, Bifone A, Gozzi A, Ragozzino D, Gross CT.** Deficient neuron-microglia signaling results in impaired functional brain connectivity and social behavior. *Nat Neurosci* 2014; 17(3):400-6
<http://www.ncbi.nlm.nih.gov/pubmed/24487234>
- **Zille M, Riabinska A, Terzi MY, Balkaya M, Prinz V, Schmerl B, Nieminen-Kelhä M, Endres M, Vajkoczy P, Pina AL.** Influence of pigment epithelium-derived factor on outcome after striatal cerebral ischemia in the mouse. *PLoS One* 2014; 9(12): e114595
<http://www.ncbi.nlm.nih.gov/pubmed/25470280>

2013

- **Amemori T, Romanyuk N, Jendelova P, Herynek V, Turnovcova K, Prochazka P, Kapcalova M, Cocks G, Price J, Sykova E.** Human conditionally immortalized neural stem cells improve locomotor function after spinal cord injury in the rat. *Stem Cell Res Ther* 2013; 4(3):68
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3706805/>
- **Dobi A, Sartori SB, Busti D, Van der Putten H, Singewald N, Shigemoto R, Ferraguti F.** Neural substrates for the distinct effects of presynaptic group III metabotropic glutamate receptors on extinction of contextual fear conditioning in mice. *Neuropharmacology* 2013; 66:274-89
<http://www.ncbi.nlm.nih.gov/pubmed/22643400>
- **Govindarajan N, Rao P, Burkhardt S, Sananbenesi F, Schlüter OM, Bradke F, Lu J, Fischer A.** Reducing HDAC6 ameliorates cognitive deficits in a mouse model for Alzheimer's disease. *EMBO Mol Med* 2013; 5(1):52-63
<http://www.ncbi.nlm.nih.gov/pubmed/23184605>

- **Greenman Y, Kuperman Y, Drori Y, Asa SL, Navon I, Forkosh O, Gil S, Stern N, Chen A.** Postnatal ablation of POMC neurons induces an obese phenotype characterized by decreased food intake and enhanced anxiety-like behavior. *Mol Endocrinol* 2013;27(7):1091-102
<http://www.ncbi.nlm.nih.gov/pubmed/23676213>
- **Grieb B, von Nicolai C, Engler G, Sharott A, Papageorgiou I, Hamel W, Engel AK, Moll CK.** Decomposition of abnormal free locomotor behavior in a rat model of Parkinson's disease. *Front Syst Neurosci* 2013; 7:(95)
<http://www.ncbi.nlm.nih.gov/pubmed/24348346>
- **Guzmán YF, Tronson NC, Jovasevic V, Sato K, Guedea AL, Mizukami H, Nishimori K, Radulovic J.** Fear-enhancing effects of septal oxytocin receptors. *Nat Neurosci* 2013; 16(9):1185-7
<http://www.ncbi.nlm.nih.gov/pubmed/23872596>
- **Hess ME, Hess S, Meyer KD, Verhagen LA, Koch L, Brönneke HS, Dietrich MO, Jordan SD, Saletore Y, Elemento O, Belgardt BF, Franz T, Horvath TL, Rüter U, Jaffrey SR, Kloppenburg P, Brüning JC.** The fat mass and obesity associated gene (*Fto*) regulates activity of the dopaminergic midbrain circuitry. *Nat Neurosci* 2013; 16(8):1042-8
<http://www.ncbi.nlm.nih.gov/pubmed/23817550>
- **Hovakimyan M, Maass F, Petersen J, Holzmann C, Witt M, Lukas J, Frech MJ, Hübner R, Rolfs A, Wree A.** Combined therapy with cyclodextrin/allopregnanolone and miglustat improves motor but not cognitive functions in Niemann-Pick Type C1 mice. *Neuroscience* 2013; 252:201-11
<http://www.ncbi.nlm.nih.gov/pubmed/23948640>
- **Jantschak F, Brosda J, Franke RT, Fink H, Möller D, Hübner H, Gmeiner P, Pertz HH.** Pharmacological profile of 2-bromoterguride at human dopamine D2, porcine serotonin 5-hydroxytryptamine 2A, and α 2C-adrenergic receptors, and its antipsychotic-like effects in rats. *J Pharmacol Exp Ther* 2013; 347(1):57-68
<http://www.ncbi.nlm.nih.gov/pubmed/23863695>
- **Karen T, Schlager GW, Bendix I, Sifringer M, Herrmann R, Pantazis C, Enot D, Keller M, Kerner T, Felderhoff-Mueser U.** Effect of propofol in the immature rat brain on short- and long-term neurodevelopmental outcome. *PLoS one* 2013; 8(5):e64480
<http://www.ncbi.nlm.nih.gov/pubmed/23737984>
- **Kerimoglu C, Agis-Balboa RC, Kranz A, Stilling R, Bahari-Javan S, Benito-Garagorri E, Halder R, Burkhardt S, Stewart AF, Fischer A.** Histone-methyltransferase MLL2 (*KMT2B*) is required for memory formation in mice. *J Neurosci* 2013; 33(8):3452-64
<http://www.ncbi.nlm.nih.gov/pubmed/23426673>
- **Kuhla A, Lange S, Holzmann C, Maass F, Petersen J, Vollmar B, Wree A.** Lifelong caloric restriction increases working memory in mice. *PLoS One* 2013; 8(7):68778
<http://www.ncbi.nlm.nih.gov/pubmed/23874758>
- **Lin Z, Shi L, Lu J, Li J, Hu H, Zuo C, Tang W, Lu Y, Bao A, Xu L.** Effects of curcumin on glucose metabolism in the brains of rats subjected to chronic unpredictable stress: a 18 F-FDG micro-PET study. *BMC Complement Altern Med* 2013; 13(1):202
<http://www.ncbi.nlm.nih.gov/pubmed/23914948>

- **Mallmann RT, Elgueta C, Sleman F, Castonguay J, Wilmes T, van den Maagdenberg A, Klugbauer N.** Ablation of CaV2.1 voltage-gated Ca(2+) channels in mouse forebrain generates multiple cognitive impairments. *PLoS One* 2013; 8(10):e78598
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3814415/>
- **Mehta NS, Wang L, Redei EE.** Sex differences in depressive, anxious behaviors and hippocampal transcript levels in a genetic rat model. *Genes Brain Behav* 2013; 12(7):695-704
<http://www.ncbi.nlm.nih.gov/pubmed/23876038>
- **Ohtsuka N, Badurek S, Busslinger M, Benes FM, Minichiello L, Rudolph U.** GABAergic neurons regulate lateral ventricular development via transcription factor Pax5. *Genesis* 2013; 51(4): 234-45
<http://www.ncbi.nlm.nih.gov/pubmed/23349049>
- **Otte DM, Barcena de Arellano ML, Bilkei-Gorzo A, Albayram O, Imbeault S, Jeung H, Alferink J, Zimmer A.** Effects of chronic D-serine elevation on animal models of depression and anxiety-related behavior. *PLoS one* 2013; 8(6):e67131
<http://www.ncbi.nlm.nih.gov/pubmed/23349049>
- **Painsipp E, Köfer MJ, Farzi A, Dischinger US, Sinner F, Herzog H, Holzer P.** Neuropeptide Y and peptide YY protect from weight loss caused by Bacille Calmette-Guérin in mice. *Br J Pharmacol* 2013; 170(5):1014-26
<http://www.ncbi.nlm.nih.gov/pubmed/23992467>
- **Pitts MW, Reeves MA, Hashimoto AC, Ogawa A, Kremer P, Seale LA, Berry MJ.** Deletion of selenoprotein M leads to obesity without cognitive deficits. *J Biol Chem* 2013; 288(36):26121-34
<http://www.ncbi.nlm.nih.gov/pubmed/23880772>
- **Reichmann F, Painsipp E, Holzer P.** Environmental enrichment and gut inflammation modify stress-induced c-Fos expression in the mouse corticolimbic system. *PLoS one* 2013; 8(1):e54811
<http://www.ncbi.nlm.nih.gov/pubmed/23349972>
- **Spinelli S, Müller T, Friedel M, Sigrist H, Lesch KP, Henkelman M, Rudin M, Seifritz E, Pryce CR.** Effects of repeated adolescent stress and serotonin transporter gene partial knockout in mice on behaviors and brain structures relevant to major depression. *Front Behav Neurosci* 2013; 7:215
<http://www.ncbi.nlm.nih.gov/pubmed/24427124>
- **Stottmann RW, Donlin M, Hafner A, Bernard A, Sinclair DA, Beier DR.** A mutation in *Tubb2b*, a human polymicrogyria gene, leads to lethality and abnormal cortical development in the mouse. *Hum Mol Genet* 2013; 22(20):4053-63
<http://www.ncbi.nlm.nih.gov/pubmed/23727838>
- **Vogel H, Montag D, Kanzleiter T, Jonas W, Matzke D, Scherneck S, Chadt A, Töle J, Kluge R, Joost HG, Schürmann A.** An Interval of the Obesity QTL *Nob3.38* within a QTL Hotspot on Chromosome 1 Modulates Behavioral Phenotypes. *PLoS One* 2013; 8(1):e53025
<http://www.ncbi.nlm.nih.gov/pubmed/23308133>
- **Wiescholleck V, Manahan-Vaughan D.** Persistent deficits in hippocampal synaptic plasticity accompany losses of hippocampus-dependent memory in a rodent model of psychosis. *Front Integr Neurosci* 2013; 7:12
<http://www.ncbi.nlm.nih.gov/pubmed/23508474>

2012

- **Andrus BM, Blizinsky K, Vedell PT, Dennis K, Shukla PK, Schaffer DJ, Radulovic J, Churchill GA, Redei EE.** Gene expression patterns in the hippocampus and amygdala of endogenous depression and chronic stress models. *Molecular Psychiatry* 2012; 17: 49-61
<http://www.ncbi.nlm.nih.gov/pubmed/21079605>
- **Bahari-Javan S, Maddalena A, Kerimoglu C, Wittnam J, Held T, Bähr M, Burkhardt S, Delalle I, Kügler S, Fischer A, Sananbenesi F.** HDAC1 regulates fear extinction in mice. *J Neurosci* 2012; 32(15): 5062-73
<http://www.ncbi.nlm.nih.gov/pubmed/22496552>
- **Freude S, Schilbach K, Hettich MM, Brönneke HS, Zemva J, Krone W, Schubert M.** Neuron-specific deletion of a single copy of the insulin-like growth factor-1 receptor gene reduces fat accumulation during aging. *Horm Metab Res* 2012; 44: 99-104
<http://www.ncbi.nlm.nih.gov/pubmed/22198814>
- **Geis C, Grünewald B, Weishaupt A, Wultsch T, Toyka KV, Reif A, Sommer C.** Human IgG directed against amphiphysin induces anxiety behavior in a rat model after intrathecal passive transfer. *J Neural Transm* 2012; 119(8): 981-5
<http://www.ncbi.nlm.nih.gov/pubmed/22331304>
- **Gruss M, Appenroth D, Flubacher A, Enzensperger C, Bock J, Fleck C, Gille G, Braun K.** 9-Methyl- β -carboline-induced cognitive enhancement is associated with elevated hippocampal dopamine levels and dendritic and synaptic proliferation. *J Neurochem* 2012; 121(6): 924-931
<http://www.ncbi.nlm.nih.gov/pubmed/22380576>
- **Guan JS, Su SC, Gao J, Joseph N, Xie Z, Zhou Y, Durak O, Zhang L, Zhu JJ, Clauser KR, Carr SA, Tsai LH.** Cdk5 is required for memory function and hippocampal plasticity via the cAMP signaling pathway. *PLoS One* 2012; 6(9): e25735
<http://www.ncbi.nlm.nih.gov/pubmed/21984943>
- **Holzmann C, Dräger D, Mix E, Hawlitschka A, Antipova V, Benecke R, Wree A.** Effects of intrastriatal botulinum neurotoxin A on the behavior of Wistar rats. *Behav Brain Res* 2012; 234(1): 107-16
<http://www.ncbi.nlm.nih.gov/pubmed/22728288>
- **Imielski Y, Schwamborn JC, Lüningschrör P, Heimann P, Holzberg M, Werner H, Leske O, Püschel AW, Memet S, Heumann R, Israel A, Kaltschmidt C, Kaltschmidt B.** Regrowing the adult brain: NF-kappaB controls functional circuit formation and tissue homeostasis in the dentate gyrus. *PLoS One* 2012; 7(2): e30838
<http://www.ncbi.nlm.nih.gov/pubmed/22312433>
- **Jawhar S, Trawicka A, Jenneckens C, Bayer TA, Wirths O.** Motor deficits, neuron loss, and reduced anxiety coinciding with axonal degeneration and intraneuronal A β aggregation in the 5XFAD mouse model of Alzheimer's disease. *Neurobiol Aging* 2012; 33(1): 19e29-40
<http://www.ncbi.nlm.nih.gov/pubmed/20619937>
- **Kastenberger I, Lutsch C, Herzog H, Schwarzer C.** Influence of sex and genetic background on anxiety-related and stress-induced behaviour of prodynorphin-deficient mice. *PLoS One* 2012; 7(3): e34251
<http://www.ncbi.nlm.nih.gov/pubmed/22479578>

- **Nakamura K, Kosugi I, Lee DY, Hafner A, Sinclair DA, Ryo A, Lu KP.** Prolyl isomerase Pin1 regulates neuronal differentiation via β -catenin. *Mol Cell Biol* 2012; 32(15):2966-78
<http://www.ncbi.nlm.nih.gov/pubmed/22645310>
- **Ribeiro PO, Valentim AM, Rodrigues P, Olsson IA, Antunes LM.** Apoptotic neurodegeneration and spatial memory are not affected by sedative and anaesthetics doses of ketamine/medetomidine combinations in adult mice. *Br J Anaesth* 2012; 108(5):807-14
<http://www.ncbi.nlm.nih.gov/pubmed/22362669>
- **Schumacher T, Krohn M, Hofrichter J, Lange C, Stenzel J, Steffen J, Dunkelmann T, Paarmann K, Fröhlich C, Uecker A, Plath AS, Sommer A, Brüning T, Heinze HJ, Pahnke J.** ABC transporters B1, C1 and G2 differentially regulate neuroregeneration in mice. *PLoS One* 2012; 7(4): e35613
<http://www.ncbi.nlm.nih.gov/pubmed/22545122>
- **Sittig LJ, Herzing LB, Xie H, Batra KK, Shukla PK, Redei EE.** Excess folate during adolescence suppresses thyroid function with permanent deficits in motivation and spatial memory. *Genes Brain Behav* 2012; 11(2):193-200
<http://www.ncbi.nlm.nih.gov/pubmed/22050771>
- **Sztainberg Y, Kuperman Y, Justice N, Chen A.** An anxiolytic role for CRF receptor type 1 in the globus pallidus. *J Neurosci* 2012; 31(48): 17416-17424
<http://www.ncbi.nlm.nih.gov/pubmed/22131403>
- **Zajda ME, Krzascik P, Hill M, Majewska MD.** Psychomotor and rewarding properties of the neurosteroids dehydroepiandrosterone sulphate and androsterone: effects on monoamine and steroid metabolism. *Acta Neurobiol Exp (Wars)* 2012; 72(1): 65-79
<http://www.ncbi.nlm.nih.gov/pubmed/22508085>

2011

- **Albayram O, Alferink J, Pitsch J, Piyanova A, Neitzert K, Poppensieker K, Mauer D, Michel K, Legler A, Becker A, Monory K, Lutz B, Zimmer A, Bilkei-Gorzo A.** Role of CB1 cannabinoid receptors on GABAergic neurons in brain aging. *Proc Natl Acad Sci* 2011; 108(27): 11256-61
<http://www.ncbi.nlm.nih.gov/pubmed/21690345>
- **Djodari-Irani A, Klein J, Banzhaf J, Joel D, Heinz A, Harnack D, Lagemann T, Juckel G, Kupsch A, Morgenstern R, Winter C.** Activity modulation of the globus pallidus and the nucleus entopeduncularis affects compulsive checking in rats. *Behav Brain Res* 2011; 219: 149-58
<http://www.ncbi.nlm.nih.gov/pubmed/21219934>
- **Engler H, Doenlen R, Engler A, Riether C, Prager G, Niemi MB, Pacheco-López G, Krügel U, Schedlowski M.** Acute amygdaloid response to systemic inflammation. *Brain Behav Immun* 2011; 25(7): 1384-92
<http://www.ncbi.nlm.nih.gov/pubmed/21521653>
- **Ermakova O, Piszczek L, Luciani L, Cavalli FMG, Ferreira T, Farley D, Rizzo S, Paolicelli RC, Al-Banchaabouchi M, Nerlov C, Moriggi R, Luscombe NM, Gross C.** Sensitized phenotypic screening identifies gene dosage sensitive region on chromosome 11 that predisposes to disease in mice. *EMBO Mol Med* 2011; 3: 50-66
<http://www.ncbi.nlm.nih.gov/pubmed/21204268>

- **Geis C, Weishaupt A, Grünewald B, Wultsch T, Reif A, Gerlach M, Dirx R, Solimena M, Perani D, Heckmann M, Toyka KV, Folli F, Sommer C.** Human stiff-person syndrome IgG induces anxious behavior in rats. *PLoS One* 2011; 6(2): e16775
<http://www.ncbi.nlm.nih.gov/pubmed/21346811>
- **Haramati S, Navon I, Issler O, Ezra-Nevo G, Gil S, Zwang R, Hornstein E, Chen A ES.** MicroRNA as repressors of stress-induced anxiety: the case of amygdalar miR-34. *J Neurosci* 2011; 31(40): 14191-14203
<http://www.ncbi.nlm.nih.gov/pubmed/21976504>
- **Kamprath K, Romo-Parra H, Häring M, Gaburro S, Döngi M, Lutz B, Pape HC.** Short-term adaption of conditioned fear responses through endocannabinoid signaling in the central amygdala. *Neuropsychopharmacology* 2011; 36(3): 652-63
<http://www.ncbi.nlm.nih.gov/pubmed/20980994>
- **Kriks S, Shim JW, Piao J, Ganat YM, Wakeman DR, Xie Z, Carrillo-Reid L, Auyeung G, Antonacci C, Buch A, Yang L, Beal MF, Surmeier DJ, Kordower JH, Tabar V, Studer L.** Dopamine neurons derived from human ES cells efficiently engraft in animal models of Parkinson's disease. *Nature* 2011; 480(7378): 547-51
<http://www.ncbi.nlm.nih.gov/pubmed/22056989>
- **Lesting J, Geiger M, Narayanan RT, Pape HC, Seidenbecher T.** Impaired extinction of fear and maintained amygdala-hippocampal theta synchrony in a mouse model of temporal lobe epilepsy. *Epilepsia* 2011; 52(2): 337-46
<http://www.ncbi.nlm.nih.gov/pubmed/21054349>
- **Liebscher I, Müller U, Teupser D, Engemaier E, Engel KMY, Ritscher L, Thor D, Sangkuhl K, Ricken A, Wurm A, Piehler D, Schmutzler S, Fuhrmann H, Albert FW, Reichenbach A, Thiery J, Schöneberg T, Schulz A.** Altered immune response in mice deficient for the G protein-coupled receptor GPR34. *J Biol Chem* 2011; 286(3): 2101-10
<http://www.ncbi.nlm.nih.gov/pubmed/21097509>
- **Mechan AO, Fowler A, Seifert N, Rieger H, Wöhrle T, Etheve S, Wyss A, Schüler G, Colletto B, Kilpert C, Aston J, Elliott JM, Goralczyk R, Mohajeri MH.** Monoamine reuptake inhibition and mood-enhancing potential of a specified oregano extract. *Br J Nutr* 2011; 105(8): 1150-63
<http://www.ncbi.nlm.nih.gov/pubmed/21205415>
- **Michalakis S, Kleppisch T, Polta SA, Wotjak CT, Koch S, Rammes G, Matt L, Becirovic E, Biel M.** Altered synaptic plasticity and behavioral abnormalities in CNGA3-deficient mice. *Genes Brain Behav* 2011; 10: 137-148
<http://www.ncbi.nlm.nih.gov/pubmed/20846178>
- **Olczak MO, Duszczak M, Mierzejewki P, Meyza K, Majewska MD.** Persistent behavioral impairments and alternations of brain dopamine system after early postnatal administration of thimerosal in rats. *Behav Brain Res* 2011; 223: 107-18
<http://www.ncbi.nlm.nih.gov/pubmed/21549155>
- **Painsipp E, Herzog H, Sperk G, Holzer P.** Sex-dependent control of murine emotional-affective behaviour in health and colitis by peptide YY and neuropeptide Y. *Br J Pharmacol* 2011; 163: 1302-14
<http://www.ncbi.nlm.nih.gov/pubmed/21410462>

- **Post AM, Weyers P, Holzer P, Painsipp E, Pauli P, Wultsch T, Reif A, Lesch KP.** Gene-environment interaction influences anxiety-like behavior in ethologically based mouse models. *Behav Brain Res* 2011; 218: 99-105
<http://www.ncbi.nlm.nih.gov/pubmed/21111005>
- **Post AM, Wultsch T, Popp S, Painsipp E, Wetzstein H, Kittel-Schneider S, Sontag TA, Lesch KP, Reif A.** The COGITAT holeboard system as a valuable tool to assess learning, memory and activity in mice. *Behav Brain Res* 2011; 220: 152-8
<http://www.ncbi.nlm.nih.gov/pubmed/21310188>
- **Sittig LJ, Shukla PK, Herzig LBK, Redei EE.** Strain-specific vulnerability to alcohol exposure in utero via hippocampal parent-of-origin expression of deiodinase-3. *FASEB J* 2011; 25: 2313-24
<http://www.ncbi.nlm.nih.gov/pubmed/21429942>
- **Xu X, Tian D, Hong X, Chen L, Xie L.** Sex-specific influence of exposure to bisphenol-A between adolescence and young adulthood on mouse behaviors. *Neuropharmacology* 2011; 61(4): 565-73
<http://www.ncbi.nlm.nih.gov/pubmed/21570416>

2010

- **Ackermann TF, KEMPE DS, Lang F, Lang UE.** Hyperactivity and enhanced curiosity of mice expressing PKB/SGK-resistant glycogen synthase kinase-3 (GSK-3). *Cell Physiol Biochem* 2010; 25: 775-86
<http://www.ncbi.nlm.nih.gov/pubmed/20511724>
- **Anier K, Malinovskaja K, Aonurm-Helm A, Zharkovsky A, Kalda A.** DNA methylation regulates cocaine-induced behavioral sensitization in mice. *Neuropsychopharmacology* 2010; 35: 2450-61
<http://www.ncbi.nlm.nih.gov/pubmed/20720536>
- **Bilkei-Gorzo A, Berner J, Zimmermann J, Wickström R, Racz I, Zimmer A.** Increased morphine analgesia and reduced side effects in mice lacking the *tac1* gene. *Br J Pharmacol* 2010; 160: 1443-52
<http://www.ncbi.nlm.nih.gov/pubmed/20590634>
- **Boy J, Schmidt T, Schumann U, Grasshoff U, Unser S, Holzmann C, Schmitt I, Karl T, Laccone F, Wolburg H, Ibrahim S, Riess O.** A transgenic mouse model of spinocerebellar ataxia type 3 resembling late disease onset and gender-specific instability of CAG repeats. *Neurobiol Dis* 2010; 37: 284-93
<http://www.ncbi.nlm.nih.gov/pubmed/19699305>
- **Gao J, Wang WY, Mao YW, Gräff J, Guan JS, Pan L, Mak G, Kim D, Su SC, Tsai LH.** A novel pathway regulates memory and plasticity via SIRT1 and miR-134. *Nature* 2010; 466(7310): 1105-9
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2928875/>
- **Klein J, Winter C, Coquery N, Heinz A, Morgenstern R, Kupsch A, Juckel G.** Lesion of the medial prefrontal cortex and the subthalamic nucleus selectively affect depression-like behavior in rats. *Behav Brain Res* 2010; 213: 73-81
<http://www.ncbi.nlm.nih.gov/pubmed/20434489>

- **Lee C, Parks GS, Civelli O.** Anxiolytic effects of the MCH1R antagonist TPI 1361-17. *J Mol Neurosci* 2010; 43(2): 132-7
<http://www.ncbi.nlm.nih.gov/pubmed/20635163>
- **Michán S, Li Y, Men-Hsiu Chou M, Parrella E, Ge H, Long JM, Allard, JS, Lewis K, Miller M, Xu W, Mervis RF, Chen J, Guerin KI, Smith LEH, McBurney MW, Sinclair DA, Baudry M, de Cabo R, Longo VD.** SIRT1 is essential for normal cognitive function and synaptic plasticity. *J Neurosci* 2010; 30(29): 9695-707
<http://www.ncbi.nlm.nih.gov/pubmed/20660252>
- **Neufeld-Cohen A, Evans AK, Getselter D, Spyroglou A, Hill A, Gil S, Tsoory M, Beuschlein F, Lowry CA, Vale W, Chen A.** Urocortin-1 and -2 double-deficient mice show robust anxiolytic phenotype and modified serotonergic activity in anxiety circuits. *Mol Psychiatry* 2010; 15(4): 426-41
<http://www.ncbi.nlm.nih.gov/pubmed/19884890>
- **Peleg S, Sananbenesi F, Zovoilis A, Burkhardt S, Bahari-Javan S, Agis-Balboa RC, Cota P, Wittnam JL, Gogol-Doering A, Opitz L, Salinas-Riester G, Dettenhofer M, Kang H, Farinelli L, Chen W, Fischer A.** Altered histone acetylation is associated with age-dependent memory impairment in mice. *Science* 2010; 328(5979): 753-6
<http://www.ncbi.nlm.nih.gov/pubmed/20448184>
- **Pilhatsch M, Winter C, Nordström K, Vennström B, Bauer M, Juckel G.** Increased depressive behaviour in mice harboring the mutant thyroid hormone receptor alpha 1. *Behav Brain Res* 2010; 214: 187-92
<http://www.ncbi.nlm.nih.gov/pubmed/20580649>
- **Regev L, Neufeld-Cohen A, Tsoory M, Kuperman Y, Getselter D, Gil S, Chen A.** Prolonged and site-specific over-expression of corticotropin-releasing factor reveals differential roles for extended amygdala nuclei in emotional regulation. *Mol Psychiatry* 2010; 16: 714-728
<http://www.ncbi.nlm.nih.gov/pubmed/20548294>
- **Sztainberg Y, Kuperman Y, Tsoory M, Lebow M, Chen A.** The anxiolytic effect of environmental enrichment is mediated via amygdala CRF receptor type 1. *Mol Psychiatry* 2010; 15(9): 905-17
<http://www.ncbi.nlm.nih.gov/pubmed/20084060>
- **Wirths O, Erck C, Martens H, Harmeier A, Geumann C, Jawhar S, Kumar S, Multhaup G, Walter J, Ingelsson M, Degerman-Gunnarsson M, Kalimo H, Huitinga I, Lannfelt L, Bayer TA.** Identification of low molecular weight pyroglutamate A β oligomers in Alzheimer disease: a novel tool for therapy and diagnosis. *J Biol Chem* 2010; 285(53): 41517-24
<http://www.ncbi.nlm.nih.gov/pubmed/20971852>

2009

- **Berna-Erro A, Braun A, Kraft R, Kleinschnitz C, Schuhmann MK, Stegner D, Wultsch T, Eilers J, Meuth SG, Stoll G, Nieswandt B.** STIM2 regulates capacitive Ca²⁺ entry in neurons and plays a key role in hypoxic neuronal cell death. *Sci Signal* 2009; 2(93): ra67
<http://www.ncbi.nlm.nih.gov/pubmed/19843959>
- **Boy J, Schmidt T, Wolburg H, Mack A, Nuber S, Böttcher M, Schmitt I, Holzmann C, Zimmermann F, Servadio A, Riess O.** Reversibility of symptoms in a conditional mouse model of spinocerebellar ataxia type 3. *Hum Mol Genet* 2009; 18(22): 4282-95

<http://www.ncbi.nlm.nih.gov/pubmed/19666958>

- **Darvas M, Morsch M, Racz I, Ahmadi S, Swandulla D, Zimmer A.** Modulation of the Ca²⁺ conductance of nicotinic acetylcholine receptors by Lypd6. *Eur Neuropsychopharmacol* 2009; 19(9): 670-81
<http://www.ncbi.nlm.nih.gov/pubmed/19403274>
- **Guan JS, Haggarty SJ, Giacometti E, Dannenberg JH, Joseph N, Gao J, Nieland TJF, Zhou Y, Wang X, Mazitschek R, Bradner JE, DePinho RA, Jaenisch R, Tsai LH.** HDAC2 negatively regulates memory formation and synaptic plasticity. *Nature* 2009; 459: 55-63
<http://www.ncbi.nlm.nih.gov/pubmed/19424149>
- **Mao R, Page DT, Merzlyak I, Kim C, Tecott LH, Janak PH, Rubenstein JLR, Sur M.** Reduced conditioned fear response in mice that lack Dlx1 and show subtype-specific loss of interneurons. *J Neurodev Disord* 2009; 1(3): 224-36
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2758250/>
- **Marques JM, Alonso I, Santos C, Silveira I, Olsson IAS.** The spatial learning phenotype of heterozygous leaner mice is robust to systematic variation of the housing environment. *Comp Med* 2009; 59(2): 129-38
<http://www.ncbi.nlm.nih.gov/pubmed/19389304>
- **Muigg P, Schreiber S, Salchner P, Bunck M, Landgraf R, Singewald N.** Differential stress-induced neuronal activation patterns in mouse lines selectively bred for high, normal or low anxiety. *PLoS One* 2009; 4(4): 1-12
<http://www.ncbi.nlm.nih.gov/pubmed/19399175>
- **Mundt A, Klein J, Joel D, Heinz A, Djodari-Irani A, Harnack D, Kupsch A, Orawa H, Juckel G, Morgenstern R, Winter C.** High-frequency stimulation of the nucleus accumbens core and shell reduces quinpirole-induced compulsive checking in rats. *Eur J Neurosci* 2009; 29: 2401-12
<http://www.ncbi.nlm.nih.gov/pubmed/19490027>
- **Nguyen NK, Sartori SB, Herzog H, Tasan R, Sperk G, Singewald N.** Effect of neuropeptide Y Y2 receptor deletion on emotional stress-induced neuronal activation in mice. *Synapse* 2009; 63(3): 236-246
<http://www.ncbi.nlm.nih.gov/pubmed/19084906>
- **Sherrin T, Todorovic C, Zeyda T, Tan CH, Hon PWT, Zhu YZ, Spiess J.** Chronic stimulation of corticotropin-releasing factor receptor 1 enhances the anxiogenic response of the cholecystokinin system. *Mol Psychiatry* 2009; 14(3): 291-307
<http://www.ncbi.nlm.nih.gov/pubmed/18195718>
- **Tasan RO, Lin S, Hetzenauer A, Singewald N, Herzog H, Sperk G.** Increased novelty-induced motor activity and reduced depression-like behavior in neuropeptide Y (NPY)-Y4 receptor knockout mice. *Neuroscience* 2009; 158: 1717-1730
<http://www.ncbi.nlm.nih.gov/pubmed/19121371>
- **Tauber SC, Ebert S, Weishaupt JH, Reich A, Nau R, Gerber J.** Stimulation of toll-like receptor 9 by chronic intraventricular unmethylated cytosine-guanine DNA infusion causes neuroinflammation and impaired spatial memory. *J Neuropathol Exp Neurol* 2009; 68(10): 1116-24
<http://www.ncbi.nlm.nih.gov/pubmed/19918123>

- **Todorovic C, Sherrin T, Pitts M, Hippel C, Rayner M, Spiess J.** Suppression of the MEK/ERK signaling pathway reverses depression-like behaviors of CRF2-deficient mice. *Neuropsychopharmacology* 2009; 34: 1416-26
<http://www.ncbi.nlm.nih.gov/pubmed/18843268>
- **Wittmann W, Schunk E, Rosskothien I, Gaburro S, Singewald N, Herzog H, Schwarzer C.** Prodynorphin-derived peptides are critical modulators of anxiety and regulate neurochemistry and corticosterone. *Neuropsychopharmacology* 2009; 34: 775-85
<http://www.ncbi.nlm.nih.gov/pubmed/18800067>
- **Wultsch T, Grimberg G, Schmitt A, Painsipp E, Wetzstein H, Breitenkamp AFS, Gründemann D, Schömig E, Lesch K-P, Gerlach M, Reif A.** Decreased anxiety in mice lacking the organic cation transporter 3. *J Neural Transm* 2009; 116: 689-97
<http://www.ncbi.nlm.nih.gov/pubmed/19280114>
- **Xu X, Ye L, Ruan Q.** Environmental enrichment induces synaptic structural modification after transient focal cerebral ischemia in rats. *Exp Biol Med* 2009; 234: 296-305
<http://www.ncbi.nlm.nih.gov/pubmed/19244205>

2008

- **Abramov U, Raud S, Innos J, Lasner H, Kurrikoff K, Tärna T, Puusaar T, Ökva K, Matsui T, Vasar E.** Different housing conditions alter the behavioural phenotype of CCK2 receptor-deficient mice. *Behav Brain Res* 2008; 193: 108-16
<http://www.ncbi.nlm.nih.gov/pubmed/18541314>
- **Appenroth D, Decker M, Tränkle C, Mohr K, Lehmann J, Fleck C.** In vivo investigation on the cholinesterase-inhibiting effects of tricyclic quinazolinimines: Scopolamine-induced cognitive impairments in rats are attenuated at low dosage and reinforced at higher dosage. *Pflugers Arch* 2009; 455: 895-901
<http://www.ncbi.nlm.nih.gov/pubmed/17717685>
- **Bilkei-Gorzo A, Otto M, Zimmer A.** Environmental modulation of anxiety-related neuronal activity and behaviors. *Behav Brain Res* 2008; 186(2): 289-92
<http://www.ncbi.nlm.nih.gov/pubmed/17904656>
- **Bilkei-Gorzo A, Rácz I, Michel K, Darvas M, Maldonado R, Zimmer A.** A common genetic predisposition to stress sensitivity and stress-induced nicotine craving. *Biol Psychiatry* 2008; 63(2): 164-71
<http://www.ncbi.nlm.nih.gov/pubmed/17570348>
- **Bourque SL, Iqbal U, Reynolds JN, Adams MA, Nakatsu K.** Perinatal iron deficiency affects locomotor behavior and water maze performance in adult male and female rats. *J Nutr* 2008; 138: 931-7
<http://www.ncbi.nlm.nih.gov/pubmed/18424604>
- **Busquet P, Hetzenauer A, Sinnegger-Brauns MJ, Striessnig J, Singewald N.** Role of L-type Ca²⁺ channel isoforms in the extinction of conditioned fear. *Learn Mem* 2008; 15: 378-86
<http://www.ncbi.nlm.nih.gov/pmc/articles/PM>
- **Grzda E, Wi[niewska RJ.** Differentiations of the effect of NMDA on the spatial learning of rats with 4 and 12 week diabetes mellitus. *Acta Neurobiol Exp* 2008; 68: 398-406
<http://www.ncbi.nlm.nih.gov/pubmed/18668163>

- **Kronenberg G, Harms C, Sobol RW, Cardozo-Pelaez F, Linhart H, Winter B, Balkaya M, Gertz K, Gay SB, Cox D, Eckart S, Ahmadi M, Juckel G, Kempermann G, Hellweg R, Sohr R, Hörtnagl H, Wilson SH, Jaenisch R, Endres M.** Folate deficiency induces neurodegeneration and brain dysfunction in mice lacking uracil DNA glycosylase. *J Neurosci* 2008; 28(28): 7219-30
<http://www.ncbi.nlm.nih.gov/pubmed/18614692>
- **Lo Iacono L, Gross C.** Alpha-Ca²⁺/calmodulin-dependent protein kinase II contributes to the developmental programming of anxiety in serotonin receptor 1A knock-out mice. *J Neurosci* 2008; 28: 6250-7
<http://www.ncbi.nlm.nih.gov/pubmed/18550767>
- **Maletínská L, Maixnerová J, Matyšková R, Haugvicová R, Pirník Z, Kiss A, Železná B.** Synergistic effect of CART (cocaine- and amphetamine-regulated transcript) peptide and cholestekinin on food intake regulation in lean mice. *BMC Neurosci* 2008; 9(101): 1-11
<http://www.ncbi.nlm.nih.gov/pmc/articles/PM>
- **Maul B, von Bohlen Und Halbach O, Becker A, Sterner-Kock A, Voigt JP, Siems WE, Grecksch G, Walther T.** Impaired spatial memory and altered dendritic spine morphology in angiotensin II type 2 receptor-deficient mice. *J Mol Med* 2008; 86(5): 563-71
<http://www.ncbi.nlm.nih.gov/pubmed/18335189>
- **Mierzejewski P, Rogowski A, Korkosz A, Bienkowski P, Filip M, Samochowiec J, Scinska A.** A protein synthesis inhibitor, cycloheximide does not alter cue-induced reinstatement of saccharin seeking. *Prog Neuropsychopharmacol Biol Psychiatry* 2008; 32(6): 1533-7
<http://www.ncbi.nlm.nih.gov/pubmed/18590943>
- **Painsipp E, Herzog H, Holzer P.** Implication of neuropeptide-Y Y2 receptors in the effects of immune stress on emotional, locomotor and social behavior of mice. *Neuropharmacology* 2008; 55(1): 117-26
<http://www.ncbi.nlm.nih.gov/pubmed/18508096>
- **Painsipp E, Wultsch T, Edelsbrunner ME, Tasan RO, Singewald N, Herzog H, Holzer P.** Reduced anxiety-like and depression-related behavior in neuropeptide Y Y4 receptor knockout mice. *Genes Brain Behav* 2008; 7(5): 532-42
<http://www.ncbi.nlm.nih.gov/pubmed/18221379>
- **Schmidt S, Richter M, Montag D, Sartorius T, Gawlik V, Hennige AM, Scherneck S, Himmelbauer H, Lutz SZ, Augustin R, Kluge R, Ruth P, Joost HG, Schürmann A.** Neuronal functions, feeding behavior, and energy balance in Slc2a3^{+/-} mice. *Am J Physiol Endocrinol Metab* 2008; 295(5): e1084-94
<http://www.ncbi.nlm.nih.gov/pubmed/18780771>
- **Winter C, Mundt A, Jalali R, Joel D, Harnack D, Morgenstern R, Juckel G, Kupsch A.** High frequency stimulation and temporary inactivation of the subthalamic nucleus reduce quinpirole-induced compulsive checking behavior in rats. *Exp Neurol* 2008; 210: 217-28
<http://www.ncbi.nlm.nih.gov/pubmed/18076877>
- **Wirths O, Breyhan H, Schäfer S, Roth C, Bayer TA.** Deficits in working memory and motor performance in the APP/PS1ki mouse model for Alzheimer's disease. *Neurobiol Aging* 2008; 29(6): 891-901
<http://www.ncbi.nlm.nih.gov/pubmed/17215062>

2007

- **Bailey A, Yoo JH, Racz I, Zimmer A, Kitchen I.** Preprodynorphin mediates locomotion and D(2) dopamine and mu-opioid receptor changes induced by chronic 'binge' cocaine administration. *J Neurochem* 2007; 102: 1817-30
<http://www.ncbi.nlm.nih.gov/pubmed/17532787>
- **Boyl PP, Di Nardo A, Mulle C, Sassoè-Pognetto, Panzanelli P, Mele A, Kneussel M, Constantini V, Perlas E, Massimi M, Vara H, Giustetto M, Witke W.** Profilin2 contributes to synaptic vesicle exocytosis, neuronal excitability, and novelty-seeking behavior. *EMBO J* 2007; 26(12): 2991-3002
<http://www.ncbi.nlm.nih.gov/pubmed/17541406>
- **Car H, Stefaniuk R, Wisniewska RJ.** Effect of MPEP in Morris water maze in adult and old rats. *Pharmacol Rep* 2007; 59: 88-93
<http://www.ncbi.nlm.nih.gov/pubmed/17377211>
- **Depino AM, Gross C.** Simultaneous assessment of autonomic function and anxiety-related behavior in BALB/c and C57BL/6 mice. *Behav Brain Res* 2007; 177: 254-60
<http://www.ncbi.nlm.nih.gov/pubmed/17204340>
- **Fischer A, Sananbenesi F, Wang X, Dobbin M, Tsai LH.** Recovery of learning and memory is associated with chromatin remodelling. *Nature* 2007; 447: 178-82
<http://www.ncbi.nlm.nih.gov/pubmed/17468743>
- **Janitzky K, Linke R, Yilmazer-Hanke DM, Grecksch G, Schwegler H.** Disrupted visceral feedback reduces locomotor activity and influences background contextual fear conditioning in C57BL/6JO1aHsd mice. *Behav Brain Res* 2007; 182: 109-18
<http://www.ncbi.nlm.nih.gov/pubmed/17586062>
- **Kowsky S, Pöppelmeyer C, Kramer ER, Falkenburger BH, Kruse A, Klein R, Schulz JB.** RET signaling does not modulate MPTP toxicity but is required for regeneration of dopaminergic axon terminals. *Proc Natl Acad Sci* 2007; 104(50): 20049-54
<http://www.ncbi.nlm.nih.gov/pubmed/18056810>
- **Painsipp E, Wultsch T, Shahbazian A, Edelsbrunner M, Kreissl MC, Schirbel A, Bock E, Pabst MA, Thoeringer CK, Huber HP, Holzer P.** Experimental gastritis in mice enhances anxiety in a gender-related manner. *Neuroscience* 2007; 150: 522-36
<http://www.ncbi.nlm.nih.gov/pubmed/17945426>
- **Papadopoulos T, Korte M, Eulenburg V, Kubota H, Retiounskaia M, Harvey RJ, Harvey K, O'Sullivan GA, Laube B, Hülsmann S, Geiger JR, Betz H.** Impaired GABAergic transmission and altered hippocampal synaptic plasticity in collybistin-deficient mice. *EMBO J* 2007; 26: 3888-99
<http://www.ncbi.nlm.nih.gov/pubmed/17690689>
- **Richter F, Hamann M, Richter A.** Chronic rotenone treatment induces behavioral effects but no pathological signs of parkinsonism in mice. *J Neurosci Res* 2007; 85: 681-91
<http://www.ncbi.nlm.nih.gov/pubmed/17171705>
- **Thoeringer CK, Wultsch T, Shahbazian A, Painsipp E, Holzer P.** Multidrug-resistance gene 1-type p-glycoprotein (MDR1 p-gp) inhibition by tariquidar impacts on neuroendocrine and behavioral processing of stress. *Psychoneuroendocrinology* 2007; 32: 1028-40
<http://www.ncbi.nlm.nih.gov/pubmed/17881135>

- **Todorovic C, Radulovic J, Jahn O, Radulovic M, Sherrin T, Hippel C, Spiess J.** Differential activation of CRF receptor subtypes removes stress-induced memory deficit and anxiety. *Eur J Neurosci* 2007; 25: 3385-97
<http://www.ncbi.nlm.nih.gov/pubmed/17553007>
- **Wicke KM, Rex A, Jongen-Relo A, Groth I, Gross G.** The guinea pig forced swim test as a new behavioral despair model to characterize potential antidepressants. *Psychopharmacology* 2007; 195: 95-102
<http://www.ncbi.nlm.nih.gov/pubmed/17646967>
- **Wultsch T, Chourbaji S, Fritzen S, Kittel S, Grunblatt E, Gerlach M, Gutknecht L, Chizat F, Golfier G, Schmitt A, Gass P, Lesch KP, Reif A.** Behavioural and expressional phenotyping of nitric oxide synthase-I knockdown animals. *J Neural Transm* 2007; 72: 69-85
<http://www.ncbi.nlm.nih.gov/pubmed/17982880>
- **Zienowicz M, Wislowska-Stanek A, Lehner M, Taracha E, Skorzewska A, Bidzinski A, Turzynska D, Sobolewska A, Walkowiak J, Maciejak P, Szyndler J, Plaznik A.** Fluoxetine attenuates the effects of pentylentetrazol on rat freezing behavior and c-Fos expression in the dorsomedial periaqueductal gray. *Neurosci Lett* 2007; 414: 252-6
<http://www.ncbi.nlm.nih.gov/pubmed/17207573>

2006

- **Baier PC, Branisa P, Koch R, Schindehutte J, Paulus W, Trenkwalder C.** Circadian distribution of motor-activity in unilaterally 6-hydroxy-dopamine lesioned rats. *Exp Brain Res* 2006; 169: 283-8
<http://www.ncbi.nlm.nih.gov/pubmed/16432693>
- **Bert B, Fink H, Hörtnagl H, Veh RW, Davies B, Theuring F, Kusserow H.** Mice over-expressing the 5-HT(1A) receptor in cortex and dentate gyrus display exaggerated locomotor and hypothermic response to 8-OH-DPAT. *Behav Brain Res* 2006; 167: 328-41
<http://www.ncbi.nlm.nih.gov/pubmed/16256213>
- **Carola V, Frazzetto G, Gross C.** Identifying interactions between genes and early environment in the mouse. *Genes Brain Behav* 2006; 5(2): 189-99
<http://www.ncbi.nlm.nih.gov/pubmed/16507009>
- **Heneka MT, Ramanathan M, Jacoby AH, Dumitrescu-Ozimek L, Bilkei-Gorzo A, Debeir T, Sastre M, Galldiks N, Zimmer A, Hoehn M, Heiss WD, Klockgether T, Staufenbiel M.** Locus ceruleus degeneration promotes alzheimer pathogenesis in amyloid precursor protein 23 transgenic mice. *J Neurosci* 2006; 26(5): 1343-54
<http://www.ncbi.nlm.nih.gov/pubmed/16452658>
- **Kuzmin A, Madjid N, Terenius L, Ogren SO, Bakalkin G.** Big dynorphin, a prodynorphin-derived peptide produces NMDA receptor-mediated effects on memory, anxiolytic-like and locomotor behavior in mice. *Neuropsychopharmacology* 2006; 31: 1928-37
<http://www.ncbi.nlm.nih.gov/pubmed/16292317>
- **Lemon N, Manahan-Vaughan D.** Dopamine D1/D5 receptors gate the acquisition of novel information through hippocampal long-term potentiation and long-term depression. *J Neurosci* 2006; 26: 7723-9
<http://www.ncbi.nlm.nih.gov/pubmed/16855100>

- **Madjid N, Tottie EE, Luttgen M, Meister B, Sandin J, Kuzmin A, Stiedl O, Ogren SO.** 5-Hydroxytryptamine 1A receptor blockade facilitates aversive learning in mice: interactions with cholinergic and glutamatergic mechanisms. *J Pharmacol Exp Ther* 2006; 316: 581-91
<http://www.ncbi.nlm.nih.gov/pubmed/16223872>
- **Meissner HO, Mrozikiewicz P, Bobkiewicz-Kozłowska T, Mscisz A, Kedzia B, Lowicka A, Reich-Bilinska H, Kapczynski W, Barchia I.** Hormone-balancing effect of pre-gelatinized organic maca (*Lepidium peruvianum Chacon*): (I) Biomedical and pharmacodynamic study on maca using clinical laboratory model on ovariectomized rats. *Int J Biomed Sci* 2006; 2(3): 260-72
<http://www.ncbi.nlm.nih.gov/pmc/articles/PM>
- **Mierzejewski P, Siemiatkowski M, Radwanska K, Szyndler J, Bienkowski P, Stefanski R, Kaczmarek L, Kostowski W.** Cycloheximide impairs acquisition but not extinction of cocaine self-administration. *Neuropharmacology* 2006; 51: 367-373
<http://www.ncbi.nlm.nih.gov/pubmed/16777145>
- **Safiulina D, Peet N, Seppet E, Zharkovsky A, Kaasik A.** Dehydroepiandrosterone inhibits complex I of the mitochondrial respiratory chain and is neurotoxic in vitro and in vivo at high concentrations. *Toxicol Sci* 2006; 93(2): 348-56
<http://www.ncbi.nlm.nih.gov/pubmed/16849397>
- **Shimshek DR, Bus T, Grinevich V, Single FN, Mack V, Sprengel R, Spergel DJ, Seeburg PH.** Impaired reproductive behavior by lack of GluR-B containing αMPA receptors but not of NMDA receptors in hypothalamic and septal neurons. *Mol Endocrinol* 2006; 20(1): 219-231
<http://www.ncbi.nlm.nih.gov/pubmed/16099814>
- **Siren AL, Radyushkin K, Boretius S, Kammer D, Riechers CC, Natt O, Sargin D, Watanabe T, Sperling S, Michaelis T, Price J, Meyer B, Frahm J, Ehrenreich H.** Global brain atrophy after unilateral parietal lesion and its prevention by erythropoietin. *Brain* 2006; 129: 480-9
<http://www.ncbi.nlm.nih.gov/pubmed/16339796>
- **Tõnissoo T, Kõks S, Meier R, Raud S, Plaas M, Vasar E, Karis A.** Heterozygous mice with Ric-8 mutation exhibit impaired spatial memory and decreased anxiety. *Behav Brain Res* 2006; 167: 42-8
<http://www.ncbi.nlm.nih.gov/pubmed/16221497>
- **Wisłowska-Stanek A, Zienowicz M, Lehner M, Taracha E, Bidzinski A, Maciejak P, Skorzewska A, Szyndler J, Plaznik A.** Midazolam inhibits neophobia-induced Fos expression in the rat hippocampus. *J Neural Transm* 2006; 113: 43-8
<http://www.ncbi.nlm.nih.gov/pubmed/16372144>

2005

- **Beglopoulos V, Montag-Sallaz M, Rohlmann A, Piechotta K, Ahmad M, Montag D, Missler M.** Neurexophilin 3 is highly localized in cortical and cerebellar regions and is functionally important for sensorimotor gating and motor coordination. *Mol Cell Biol* 2005; 25(16): 7278-88
<http://www.ncbi.nlm.nih.gov/pubmed/16055736>
- **Beraki S, Aronsson F, Karlsson H, Ogren SO, Kristensson K.** Influenza A virus infection causes alterations in expression of synaptic regulatory genes combined with changes in cognitive and emotional behaviors in mice. *Mol Psychiatry* 2005; 10(3): 299-308

<http://www.ncbi.nlm.nih.gov/pubmed/15241434>

- **Bert B, Dere E, Wilhelmi N, Kusserow H, Theuring F, Huston JP, Fink H.** Transient overexpression of the 5-HT1A receptor impairs water-maze but not hole-board performance. *Neurobiol Learn Mem* 2005; 84(1): 57-68
<http://www.ncbi.nlm.nih.gov/pubmed/15936683>
- **Fischer A, Sananbenesi F, Pang PT, Tsai L.** Opposing roles of transient and prolonged expression of p25 in synaptic plasticity and hippocampus-dependent memory. *Neuron* 2005; 48: 825-38
<http://www.ncbi.nlm.nih.gov/pubmed/16337919>
- **Kuteeva E, Hökfelt T, Ogren SO.** Behavioural characterisation of transgenic mice overexpressing galanin under the PDGF-B promoter. *Neuropeptides* 2005; 39: 299-304
<http://www.ncbi.nlm.nih.gov/pubmed/15944026>
- **Kuteeva E, Hökfelt T, Ogren SO.** Behavioural characterisation of young adult transgenic mice overexpressing galanin under the PDGF-B promoter. *Regul Pept* 2005; 125: 67-78
<http://www.ncbi.nlm.nih.gov/pubmed/15582716>
- **Langen B, Egerland U, Bernoster K, Dost R, Unverferth K, Rundfeldt C.** Characterization in rats of the anxiolytic potential of ELB139 [1-(4-chlorophenyl)-4-piperidin-1-yl-1,5-dihydroimidazol-2-on], a new agonist at the benzodiazepine binding site of the GABAA receptor. *J Pharmacol Exp Ther* 2005; 314(2): 717-24
<http://www.ncbi.nlm.nih.gov/pubmed/15860576>
- **Park SK, Nguyen MD, Fischer A, Luke MP, Affar el B, Dieffenbach PB, Tseng HC, Shi Y, Tsai LH.** Par-4 links dopamine signaling and depression. *Cell* 2005; 122: 275-87
<http://www.ncbi.nlm.nih.gov/pubmed/16051151>
- **Rex A, Voigt JP, Fink H.** Anxiety but not arousal increases 5-hydroxytryptamine release in the rat ventral hippocampus in vivo. *Eur J Neurosci* 2005; 22(5): 1185-9
<http://www.ncbi.nlm.nih.gov/pubmed/16176361>
- **Rusu AS, Krackow S.** Agonistic onset marks emotional changes and dispersal propensity in wild house mouse males (*Mus domesticus*). *J Comp Psychol* 2005; 119(1): 58-66
<http://www.ncbi.nlm.nih.gov/pubmed/15740430>
- **Strekalova T, Spanagel R, Dolgov O, Bartsch D.** Stress-induced hyperlocomotion as a confounding factor in anxiety and depression models in mice. *Behav Pharmacol* 2005; 16: 171-80.
<http://www.ncbi.nlm.nih.gov/pubmed/15864072>
- **Winter B, Juckel G, Viktorov I, Katchanov J, Gietz A, Sohr R, Balkaya M, Hortnagl H, Endres M.** Anxious and hyperactive phenotype following brief ischemic episodes in mice. *Biol Psychiatry* 2005; 57: 1166-75
<http://www.ncbi.nlm.nih.gov/pubmed/15866557>

2004

- **Bilkei-Gorzo A, Racz I, Michel K, Zimmer A, Klingmuller D, Zimmer A.** Behavioral phenotype of pre-proenkephalin-deficient mice on diverse congenic backgrounds. *Psychopharmacology* 2004; 176: 343-52
<http://www.ncbi.nlm.nih.gov/pubmed/15197532>

- **Ebner K, Rupniak NM, Saria A, Singewald N.** Substance P in the medial amygdala: emotional stress-sensitive release and modulation of anxiety-related behavior in rats. *Proc Natl Acad Sci USA* 2004; 101: 4280-5
<http://www.ncbi.nlm.nih.gov/pubmed/15024126>
- **Gerber J, Böttcher T, Hahn M, Siemer A, Bunkowski S, Nau R.** Increased mortality and spatial memory deficits in TNF-alpha-deficient mice in ceftriaxone-treated experimental pneumococcal meningitis. *Neurobiol Dis* 2004; 16: 133-8
<http://www.ncbi.nlm.nih.gov/pubmed/15207270>
- **Kittner H, Hoffmann E, Krügel U, Illes P.** P2 receptor-mediated effects on the open field behaviour of rats in comparison with behavioural responses induced by the stimulation of dopamine D2-like and by the blockade of ionotropic glutamate receptors. *Behav Brain Res* 2004; 149: 197-208
<http://www.ncbi.nlm.nih.gov/pubmed/15129782>
- **Kusserow H, Davies B, Hörtnagl H, Voigt I, Stroh T, Bert B, Deng DR, Fink H, Veh RW, Theuring F.** Reduced anxiety-related behaviour in transgenic mice overexpressing serotonin 1A receptors. *Brain Res Mol Brain Res* 2004; 129: 104-16
<http://www.ncbi.nlm.nih.gov/pubmed/15469887>
- **Rex A, Spsychalla M, Fink H.** Treatment with reduced nicotinamide adenine dinucleotide (NADH) improves water maze performance in old Wistar rats. *Behav Brain Res* 2004; 154: 149-53
<http://www.ncbi.nlm.nih.gov/pubmed/15302120>
- **Rex A, Voigt JP, Gustedt C, Beckett S, Fink H.** Anxiolytic-like profile in Wistar, but not Sprague-Dawley rats in the social interaction test. *Psychopharmacology* 2004; 177: 23-34
<http://www.ncbi.nlm.nih.gov/pubmed/15167980>
- **Tezval H, Jahn O, Todorovic C, Sasse A, Eckart K, Spiess J.** Cortagine, a specific agonist of corticotropin-releasing factor receptor subtype 1, is anxiogenic and antidepressive in the mouse model. *Proc Natl Acad Sci* 2004; 101(25): 9468-73
<http://www.ncbi.nlm.nih.gov/pmc/articles/PM>
- **Werner C, Raivich G, Cowen M, Strekalova T, Sillaber I, Buters JT, Spanagel R, Hofmann F.** Importance of NO/cGMP signalling via cGMP-dependent protein kinase II for controlling emotionality and neurobehavioural effects of alcohol. *Eur J Neurosci* 2004; 20: 3498-506
<http://www.ncbi.nlm.nih.gov/pubmed/15610182>
- **Winter B, Bert B, Fink H, Dirnagl U, Endres M.** Dysexecutive syndrome after mild cerebral ischemia? Mice learn normally but have deficits in strategy switching. *Stroke* 2004; 35: 191-5
<http://www.ncbi.nlm.nih.gov/pubmed/14671235>

2003

- **Franke H, Kittner H, Grosche J, Illes P.** Enhanced P2Y1 receptor expression in the brain after sensitisation with d-amphetamine. *Psychopharmacology* 2003; 167: 187-94
<http://www.ncbi.nlm.nih.gov/pubmed/12652343>
- **Kiefer F, Jahn H, Koester A, Montkowski A, Reinscheid RK, Wiedemann K.** Involvement of NMDA receptors in alcohol-mediated behavior: mice with reduced affinity of the NMDA R1 glycine binding site display an attenuated sensitivity to ethanol. *Biol Psychiatry* 2003; 53: 345-51

<http://www.ncbi.nlm.nih.gov/pubmed/12586454>

- **Kuzmin A, Sandin J, Terenius L, Ogren SO.** Acquisition, expression, and reinstatement of ethanol-induced conditioned place preference in mice: effects of opioid receptor-like 1 receptor agonists and naloxone. *J Pharmacol Exp Ther* 2003; 304(1): 310-8
<http://www.ncbi.nlm.nih.gov/pubmed/12490606>
- **Montag-Sallaz M, Montag D.** Severe cognitive and motor coordination deficits in Tenascin-R-deficient mice. *Genes Brain Behav* 2003; 2: 20-31
<http://www.ncbi.nlm.nih.gov/pubmed/12882316>
- **Muller MB, Zimmermann S, Sillaber I, Hagemeyer TP, Deussing JM, Timpl P, Kormann MS, Droste SK, Kuhn R, Reul JM, Holsboer F, Wurst W.** Limbic corticotropin-releasing hormone receptor 1 mediates anxiety-related behavior and hormonal adaptation to stress. *Nat Neurosci* 2003; 6: 1100-7
<http://www.ncbi.nlm.nih.gov/pubmed/12973355>
- **Racz I, Bilkei-Gorzo A, Toth ZE, Michel K, Palkovits M, Zimmer A.** A critical role for the cannabinoid CB1 receptors in alcohol dependence and stress-stimulated ethanol drinking. *J Neurosci* 2003; 23(6): 2453-8
<http://www.ncbi.nlm.nih.gov/pubmed/12657705>
- **Sienkiewicz-Jarosz H, Czlonkowska AI, Lehner M, Maciejak P, Siemiakowski M, Szyndler J, Wislowska A, Zienowicz M, Bidzinski A, Kostowski W, Plaznik A.** Lack of changes in cortical [3H]-muscimol binding in rats sensitized to nicotine-induced enhancement of dopamine metabolism. *Pol J Pharmacol* 2003; 55(2): 165-70
<http://www.ncbi.nlm.nih.gov/pubmed/12926543>
- **Sienkiewicz-Jarosz H, Maciejak P, Bidzinski A, Szyndler J, Siemiakowski M, Czlonkowska AI, Lehner M, Plaznik A.** Exploratory activity and a conditioned fear response: correlation with cortical and subcortical binding of the alpha4beta2 nicotinic receptor agonist [3H]-Epibatidine. *Pol J Pharmacol* 2003; 55: 17-23
<http://www.ncbi.nlm.nih.gov/pubmed/12856821>
- **Sienkiewicz-Jarosz H, Maciejak P, Krzascik P, Czlonkowska AI, Szyndler J, Bidzinski A, Kostowski W, Plaznik A.** The effects of central administration of physostigmine in two models of anxiety. *Pharmacol Biochem Behav* 2003; 75: 491-496
<http://www.ncbi.nlm.nih.gov/pubmed/12873642>
- **Sienkiewicz-Jarosz H, Szyndler J, Czlonkowska AI, Siemiakowski M, Maciejak P, Wislowska A, Zienowicz M, Lehner M, Turzynska D, Bidzinski A, Plaznik A.** Rat behavior in two models of anxiety and brain (3H)muscimol binding: pharmacological, correlation and multifactor analysis. *Behav Brain Res* 2003; 145: 17-22
<http://www.ncbi.nlm.nih.gov/pubmed/14529801>

2002

- **Bert B, Fink H, Huston JP, Voits M.** Fischer 344 and Wistar rats differ in anxiety and habituation but not in water maze performance. *Neurobiol Learn Mem* 2002; 78: 11-22
<http://www.ncbi.nlm.nih.gov/pubmed/12071664>
- **Bilkei-Gorzo A, Racz I, Michel K, Zimmer A.** Diminished anxiety- and depression-related behaviors in mice with selective deletion of the Tac1 gene. *J Neurosci* 2002; 22(22): 10046-52
<http://www.ncbi.nlm.nih.gov/pubmed/12427862>

- **Brakebusch C, Seidenbecher C, Asztely F, Rauch U, Matthies H, Meyer H, Krug M, Bockers T, Zhou X, Kreutz M, Montag D, Gundelfinger E, Fassler R.** Brevican-deficient mice display impaired hippocampal CA1 long-term potentiation but show no obvious deficits in learning and memory. *Mol Cell Biol* 2002; 21: 7417-27
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC135663/>
- **Montag-Sallaz M, Schachner M, Montag D.** Misguided axonal projections, neural cell adhesion molecule 180 mRNA upregulation, and altered behavior in mice deficient for the close homolog of L1. *Mol Cell Biol* 2002; 22: 7967-81
<http://www.ncbi.nlm.nih.gov/pubmed/12391163>
- **Van Gaalen MM, Stenzel-Poore M, Holsboer F, Steckler T.** Effects of transgenic overproduction of CRH on anxiety-like behaviour. *Eur J Neurosci* 2002; 15: 2007-15
<http://www.ncbi.nlm.nih.gov/pubmed/12099906>

2001

- **Azad SC, Marsicano G, Eberlein I, Putzke J, Zieglgansberger W, Spanagel R, Lutz B.** Differential role of the nitric oxide pathway on delta-(9)-THC-induced central nervous system effects in the mouse. *Eur J Neurosci* 2001; 13: 561-8
<http://www.ncbi.nlm.nih.gov/pubmed/11168564>
- **Eckart K, Jahn O, Radulovic J, Tezval H, Van Werven L, Spiess J.** A single amino acid serves as an affinity switch between the receptor and the binding protein of corticotropin-releasing factor: Implications for the design of agonists and antagonist. *Proc Natl Acad Sci USA* 2001; 98(20): 11142-7
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC58697/>
- **Keck ME, Welt T, Wigger A, Renner U, Engelmann M, Holsboer F, Landgraf R.** The anxiolytic effect of the CRH(1) receptor antagonist R121919 depends on innate emotionality in rats. *Eur J Neurosci* 2001; 13: 373-380
<http://www.ncbi.nlm.nih.gov/pubmed/11168542>
- **Kruegel U, Kittner H, Franke H, Illes P.** Stimulation of P2 receptors in the ventral tegmental area enhances dopaminergic mechanisms in vivo. *Neuropharmacology* 2001; 40: 1084-93
<http://www.ncbi.nlm.nih.gov/pubmed/11406200>
- **Siemiakowski M, Maciejak P, Sienkiewicz-Jarosz H, Czlonkowska AI, Szyndler J, Gryczynska A, Plaznik A.** Opposite effects of olanzapine and haloperidol in rat ultrasonic vocalization test. *Pol J Pharmacol* 2001; 53(6): 669-73
<http://www.ncbi.nlm.nih.gov/pubmed/11985344>
- **Stobrawa S, Breiderhoff T, Takamori S, Engel D, Schweizer M, Zdebik A, Boesl M, Ruether K, Jahn H, Draguhn A, Jahn R, Jentsch T.** Disruption of CIC-3, a chloride channel expressed on synaptic vesicles, leads to a loss of the hippocampus. *Neuron* 2001; 29: 185-96
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4146203/>

2000

- **Keck ME, Hatzinger M, Wotjak CT, Landgraf R, Holsboer F, Neumann ID.** Ageing alters intrahypothalamic release patterns of vasopressin and oxytocin in rats. *Eur J Neurosci* 2000; 12: 1487-94
<http://www.ncbi.nlm.nih.gov/pubmed/10762376>
- **Kew JNC, Koester A, Moreau JL, Jenck F, Ouagazzal AM, Mutel V, Richards JG, Trube G, Fischer G, Montkowski A, Hundt W, Reinscheid RK, Pauly-Evers M, Kemp JA, Bluethmann H.** Functional consequences of reduction in NMDA receptor glycine affinity in mice carrying targeted point mutations in the glycine binding site. *J Neurosci* 2000; 20(11): 4037-49
<http://www.ncbi.nlm.nih.gov/pubmed/10818139>
- **Kishimoto T, Radulovic J, Radulovic M, Lin C, Schrick C, Hooshmand F, Hermanson O, Rosenfeld M, Spiess J.** Deletion of Crhr2 reveals an anxiolytic role for corticotropin-releasing hormone receptor-2. *Nat Genet* 2000; 24: 415-9
<http://www.ncbi.nlm.nih.gov/pubmed/10742109>
- **Kittner H, Krügel U, Hoffmann E, Illes P.** Effects of intra-accumbens injection of 2-methylthio ATP: a combined open field and electroencephalographic study in rats. *Psychopharmacology* 2000; 150: 123-31
<http://www.ncbi.nlm.nih.gov/pubmed/10907665>
- **Sauvage M, Brabet P, Holsboer F, Bockaert J, Steckler T.** Mild deficits in mice lacking pituitary adenylate cyclase-activating polypeptide receptor type 1 (PAC1) performing on memory tasks. *Brain Res Mol Brain Res* 2000; 84: 79-89
<http://www.ncbi.nlm.nih.gov/pubmed/11113534>
- **Siemiakowski M, Rokicki D, Czlonkowska AI, Sienkiewicz-Jarosz H, Bidzinski A, Plaznik A.** Locomotor activity and a conditioned fear response: correlation with cortical and subcortical binding of the dopamine D1 receptor antagonist. *Neuroreport* 2000; 11(18): 3953-6
<http://www.ncbi.nlm.nih.gov/pubmed/11192608>
- **Wellmer A, Noeske C, Gerber J, Munzel U, Nau R.** Spatial memory and learning deficits after experimental pneumococcal meningitis in mice. *Neurosci Lett* 2000; 296: 137-40
<http://www.ncbi.nlm.nih.gov/pubmed/11109000>

1999

- **Koester A, Montkowski A, Schulz S, Stuebe EM, Knaudt K, Jenck F, Moreau JL, Nothacker HP, Civelli O, Reinscheid RK.** Targeted disruption of the orphanin FQ/nociceptin gene increases stress susceptibility and impairs stress adaptation in mice. *Proc Natl Acad Sci* 1999; 96: 10444-9
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC17908/>
- **Rupprecht R, Koch M, Montkowski A, Lancel M, Faulhaber J, Harting J, Spanagel R.** Assessment of neuroleptic-like properties of progesterone. *Psychopharmacology* 1999; 143: 29-38
<http://www.ncbi.nlm.nih.gov/pubmed/10227077>

1998

- **Radulovic J, Kammermeier J, Spiess J.** Generalization of fear responses in C57BL/6N mice subjected to one-trial foreground contextual fear conditioning. *Behav Brain Res* 1998; 95: 179-89
<http://www.ncbi.nlm.nih.gov/pubmed/9806438>
- **Schroeder U, Schroeder H, Darius J, Grecksch G, Sabel BA.** Simulation of psychosis by continuous delivery of phencyclidine from controlled-release polymer implants. *Behav Brain Res* 1998; 97: 59-68
<http://www.ncbi.nlm.nih.gov/pubmed/9867231>
- **Sillaber I, Montkowski A, Landgraf R, Barden N, Holsboer F, Spanagel R.** Enhanced morphine-induced behavioural effects and dopamine release in the nucleus accumbens in a transgenic mouse model of impaired glucocorticoid (type II) receptor function: influence of long-term treatment with the antidepressant Moclobemide. *Neuroscience* 1998; 85(2):415-25
<http://www.ncbi.nlm.nih.gov/pubmed/9622241>
- **Strohle A, Poettig M, Barden N, Holsboer F, Montkowski, A.** Age- and stimulus-dependent changes in anxiety-related behaviour of transgenic mice with GR dysfunction. *Neuroreport* 1998; 9: 2099-102
<http://www.ncbi.nlm.nih.gov/pubmed/9674601>
- **Timpl P, Spanagel R, Sillaber I, Kresse A, Reul JM, Stalla GK, Blanquet V, Steckler T, Holsboer F, Wurst W.** Impaired stress response and reduced anxiety in mice lacking a functional corticotropin-releasing hormone receptor 1. *Nat Genet* 1998; 19: 162-6
<http://www.ncbi.nlm.nih.gov/pubmed/9620773>

1997

- **Montkowski A, Poettig M, Mederer A, Holsboer F.** Behavioural performance in three substrains of mouse strain 129. *Brain Res* 1997; 762: 12-8
<http://www.ncbi.nlm.nih.gov/pubmed/9262153>
- **Skutella T, Schwarting RK, Huston JP, Sillaber I, Probst JC, Holsboer F, Spanagel R.** Infusions of tyrosine hydroxylase antisense oligodeoxynucleotide into substantia nigra of the rat: effects on tyrosine hydroxylase mRNA and protein content, striatal dopamine release and behaviour. *Eur J Neurosci* 1997; 9: 210-20
<http://www.ncbi.nlm.nih.gov/pubmed/9058042>
- **Strohle A, Jahn H, Montkowski A, Liebsch G, Boll E, Landgraf R, Holsboer F, Wiedemann K.** Central and peripheral administration of atriopeptin is anxiolytic in rats. *Neuroendocrinology* 1997; 65: 210-5
<http://www.ncbi.nlm.nih.gov/pubmed/9088002>

1996

- **Shoab M, Shippenberg TS.** Adrenalectomy attenuates nicotine-induced dopamine release and locomotor activity in rats. *Psychopharmacology* 1996; 128: 343-50



<http://www.ncbi.nlm.nih.gov/pubmed/8986004>

- **Spanagel B, Stöhr T, Barden N, Holsboer F.** Morphine-induced locomotor and neurochemical stimulation is enhanced in transgenic mice with impaired glucocorticoid receptor function. *J Neuroendocrinol* 1996; 8: 93-7
<http://www.ncbi.nlm.nih.gov/pubmed/8868255>

1995

- **Shoaib M, Spanagel R, Stohr T, Shippenberg TS.** Strain differences in the rewarding and dopamine-releasing effects of morphine in rats. *Psychopharmacology* 1995; 117(2): 240-7
<http://www.ncbi.nlm.nih.gov/pubmed/7753973>



Service & Warranty

TSE Systems offers a Two (2) Years ALL-IN Premium Warranty with all new products, including:

- 24/7 technical hotline
- Remote maintenance and update function
- On-site visits upon necessity
- Free replacement parts during warranty

After the expiry of the warranty period, TSE Systems offers comprehensive extensions of the warranty or economical maintenance and repair contracts to ensure the continued smooth running of your instruments. Please contact us for further details.